



## Effects of salinity and irrigation water management on soil and tomato in drip irrigation

M. Noshadi\*, S. Fahandej, A.R. Sepaskhah

*Department of Irrigation Engineering, Shiraz University, Iran.*

\*Corresponding author. E-mail: noshadi@shirazu.ac.ir

Received 26 July 2012; Accepted after revision 9 December 2012; Published online 1 February 2013

---

### Abstract

In this research, the effects of irrigation with saline and fresh water through drip irrigation method and using two irrigation management strategies:  $M_1$ ,  $M_2$  ( $M_1$  is irrigation with fresh water in alternative with saline water,  $M_2$  is saline water in first half time of each irrigation event and fresh water in the second half), on the yield, water productivity (WP), soil salinity, plant height and diameter, fruit water content, fruit density, "L", "a" and "b" colorimetric factors, textures, strength, sodium, calcium and nitrogen concentrations were investigated. The experiments were conducted in a randomized completely block design as split plot with three replications in which management strategies were the main plots and subplots were different levels of salinity, 0.68, 2, 4, 6 and 8 dS/m ( $S_0$ ,  $S_1$ ,  $S_2$ ,  $S_3$ ,  $S_4$ , respectively). Highest and lowest yields were in  $M_1S_1$  (59.12 t/ha) and  $M_2S_4$  (18.81 t/ha) treatments, respectively. The yield difference between  $M_1$  and  $M_2$  management strategies were significant at 5% level of probability and the average yield of  $M_2$  was 24.20% less than the  $M_1$  treatment. Applied irrigation water was decreased with increasing salinity levels because leaf area and leaf transpiration was reduced. Therefore, water productivity was increased, so that the highest water productivity was in the  $M_1S_4$  treatment. To evaluate the use of saline water on soil,  $EC_e$  was measured in each plot at four layers in soil. Highest  $EC_e$  was in the  $S_4$  salinity level in both management strategies. Based on soil salinity and crop yield,  $M_1$  and  $M_2$  management strategies were suitable at lower levels of salinities (0.68 and 2 dS/m) and salinities over 4 dS/m, respectively. Furthermore,  $M_1$  management strategy, due to more efficient leaching in the surface layers of soil, was more appropriate than  $M_2$  management strategy.

**Keywords:** Salinity; Irrigation management; Tomato; Water productivity.

---

## **Introduction**

Because of increasing world population and thereafter increasing demand for food, use of fresh water resources has increased (Wallace, 2000). On the other hand, the world's fresh water resources are limited; that forced farmers to use low quality waters. About 12% of Iran's surface waters are saline; so the role of saline and brackish waters in the future would be undeniable (Anonymous, 2009).

In general, water quality has significant effect on yield and application of saline water has reduced the amount of yield (Murtaza et al., 2006). Appropriate irrigation methods and irrigation management strategies can reduce the effects of salinity of irrigation water on soil and crop and increase water productivity. Pasternak et al. (1986) used drip irrigation system with brackish ( $EC_i=6.2$  dS/m) and fresh ( $EC_i=1.2$  dS/m) waters over a period of three growing seasons to were about 44% less than fresh water. Field experiments were conducted on a sandy loam soil by Narsh et al. (1993) to evaluate the changes in soil water and salinity when conjunctive irrigation method with fresh (0.6 dS/m) and saline (12 dS/m) waters were applied in various cyclic/mixing modes. Relative yield with saline water was reduced to 60%. The yield in cyclic irrigations with fresh and saline waters was 7-11% more than mixing method in equal proportions. It was concluded when conjunctive use of fresh and saline waters were applied for the production of wheat, water productivity was higher than cyclic use of fresh and saline waters, when fresh water was applied at the initial stages (pre-irrigation and/first post-sowing irrigation) and saline water was applied at the later growth periods when it can tolerate the salts. Pasternak (1995) studied the tomato irrigation management with saline water. Results showed that when tomatoes were irrigated with saline water (7.5 dS/m), the total yield was reduced by 60% relative to the control. However, when irrigation with saline water started at the appearance of the fourth or the eleventh leaf, a water salinity of 7.5 dS/m reduced the yield by only about 30%. Malash et al. (2005) studied the effect of two water management strategies i.e. alternate and mixed supply of fresh water and saline water in six ratios applied through drip and furrow method on tomato yield and growth. Results showed that the highest yield obtained (3.2 Kg/plant) was the result of the combination of drip system and mixed management practice using a ratio of 60% fresh water with 4% saline water. Effects of drip irrigation using saline water on the production of tomato, in silt loam soil texture and

semi-humid climates in China were investigated by Wan et al. (2010). Results showed that salinity in the range of 1.1-4.9 dS/m had little impact on yield and had the greatest impact on the cumulative amount of water used by plant and water productivity. Therefore, with increasing salinity, water productivity increased. Malash et al. (2008) studied there sponse of tomato to irrigation with saline water applied by different irrigation methods and water management strategies. The results indicated that salinity (at 3 dS/m and above) significantly reduced leaf area, height and dry weight of plant as well as fruit weight and number and hence total yield, but increased fruit T.S.S. content. Water use efficiency (WUE) was increased by using water with low and moderate salinity levels (2 and 3 dS/m) as compared to those obtained with non-saline water (0.55 dS/m) or the highest salinity level (4.5 dS/m). Salinity increased Na, Cl and Mg contents as well as dry matter percentage, but decreased N, P, K and Ca contents in leaves of plants. Drip irrigation enhanced tomato growth, yield and WUE under both saline and non-saline conditions, but showed more advantages under saline conditions as compared with furrow irrigation. Drip irrigation method did not allow salt accumulation in root zone (wetted area beneath the emitters and the plants). Using saline water up to 3 dS/m produced yield that was not significantly differ than that produced by non saline water if applied by drip irrigation and blended water management.

In this research, effects of saline and fresh water in drip irrigation under two irrigation management strategies  $M_1$ ,  $M_2$  ( $M_1$  is alternative irrigation with fresh water and saline water,  $M_2$  is saline water in first half of each irrigation event and fresh water in the second half), on the yield and quality parameters of tomato, water productivity (WP), soil salinity, plant height and diameter, fruit water content, fruit density, "L", "a" and "b" colorimetric factors, textures, strength, sodium, calcium and nitrogen concentrations were investigated.

## **Materials and Methods**

This research was conducted in the College of Agriculture Shiraz University, located 16 km north of Shiraz (latitude  $36^{\circ} 29'$ , longitude  $32^{\circ} 52'$  and altitude 1810 m) in 2010. Soil physical characteristics and irrigation water analysis are shown in Tables 1 and 2, respectively. The field area was  $319 \text{ m}^2$  including 30 plots with dimensions  $2 \times 3 \text{ m}$  with three replications. To prevent water seepage from a plot to adjacent plots, 1 m spacing was

considered between plots (Figure 1). Experimental design was split plot, in which, water management strategy was the main plots ( $M_1$ , alternative irrigation with fresh water and saline water;  $M_2$ , is saline water in first half time of each irrigation event and fresh water in the second half) and subplots were four salinity levels of irrigation water i.e. 0.68 (control), 2, 4, 6, 8 dS/m as  $S_0, S_1, S_2, S_3, S_4$ , respectively. The soil moisture was measured by neutron probe and irrigation interval was 3 days. All treatments were irrigated with fresh water until the appearance of the fourth or the eleventh leaf and then saline irrigation water treatments were applied. For preparing the saline waters two reservoirs (one for saline water and another for fresh water) was considered. The saline water was performed by NaCl and  $CaCl_2$ . The irrigation water was mixed of saline and fresh water until the desired salinity was obtained. For the control of this mixing the valves were considered and the  $E_c$  of mixed water were measured by  $E_c$ -meter before each irrigation. The variation of  $E_c$  was  $\pm 5\%$ . Irrigation water was applied by surface drip irrigation. Tomatoes were Transplanting on the 13<sup>th</sup> of April, at 3 rows in each plot, with a distance of 60 cm between rows. The amount of soil moisture was measured by the Neutron probe. Irrigation depth was determined according to following equation:




Figure 1. Experimental design.

Table 1. Soil physical properties in different soil depth.

Depth (cm)	Soil texture	Particles(%)			$\rho_b(g.cm^{-3})^1$	$PWP(cm^3.cm^{-3})^2$	$FC(cm^3.cm^{-3})^3$	$EC(ds/m)^4$	pH	OC% <sup>5</sup>
		Clay	silt	sand						
0-30	Silty clay loam	55	35	10	1.23	0.12	0.27	0.686	7.68	0.844
30-60	Silty loam	49	12	39	1.46	0.16	0.31	0.476	7.54	0.591
60-90	Silty loam	60	11	29	1.46	0.16	0.31	0.471	7.41	0.493
90-120	Silty loam	63	11	26	1.46	0.16	0.31	0.450	7.4	0.410

1- Bulk density, 2- Permanent wilting point, 3- Field capacity, 4- Soil electrical conductivity, 5- Organic carbon.

Table 2. Chemical analysis of fresh irrigation water.

Quantity	Parameters
7.65	pH
0.679	EC (dS/m)
1.7	Chloride (meq/l)
2.9	Calcium (meq/l)
5.2	Magnesium (meq/l)
0.76	Sodium (meq/l)
0.015	Potassium (meq/l)
7.4	Bicarbonate (meq/l)

$$d = \frac{(Fc - \theta_v) \times R_z}{100} \quad (1)$$

Where:  $d$  is the irrigation water depth (cm);  $\theta_v$  is the volumetric soil water content within the root depth before irrigation (%);  $Fc$  is the volumetric soil water content at field capacity (%);  $R_z$  is the root depth (cm). In Eq. (1)  $R_z$  is varied with time and obtained from the following equation (Borg and Grimes, 1986):

$$Z_r = RD_m [0.5 + 0.5 \sin (3.03 DA_s / DT_m - 1.47)] \quad (2)$$

Where:  $Z_r$  is the root depth of the given day;  $DA_s$  is the number of days after planting,  $DT_m$  is the number of days to reach the maximum root depth and  $RD_m$  is the maximum depth of plant roots. Values of  $DT_m$  and  $RD_m$ , were considered 80 days and 90 cm, respectively. Nitrogen was applied with a rate of 150 kg ha<sup>-1</sup> urea through irrigation water (70 kg ha<sup>-1</sup> at 5 June 2010 and 80 kg ha<sup>-1</sup> at 22 June, 2010). Tomatoes yield were harvested from middle row of each plot at two times (6 and 26 September, 2010) and fruits weight were determined. Furthermore, plant height and canopy diameter were measured. At the end of the growing season, after the last irrigation with saline water for both irrigation management strategies, soil was sampled in each plot from 0-30, 30-50, 50-70 and 70-90 cm depths and for each layer, electrical conductivity of saturated extract ( $EC_e$ ) was determined. In order to evaluate the tomatoes quality, fruit water content, fruit density, colorimetric characteristics, tomato texture strength and the amount of sodium, calcium and total nitrogen concentration were measured.

### Measurement of plant parameters

#### Plant height and canopy diameter

To evaluate the effect of treatments on plant height and canopy diameter, when the vegetative growth stopped, these parameter were measured in three randomly selected plants from middle row of each plot (Table 3).

#### Fruit water content

On average 93% of tomato weight is water (Holland et al., 1991). In higher water content, the tomato texture strength and durability reduce and its transportation is difficult. To determine the water content of fruits, three medium sized tomatoes per plot were selected randomly and the fresh weight was measured and then dried inside oven (65 °C) in paper envelope. After drying, fruit weights were measured again. Fruit water content obtained from the following equation:

$$w = \frac{w_1 - w_2}{w_1} \times 100 \quad (3)$$

Where:  $w$  is the water content (%),  $w_1$  is the fresh fruit weight (gr) and  $w_2$  is the dried fruit weight (gr).

#### Fruit density

One common use of tomato is in paste and sauce production. The fruits with higher density are more suitable for these purposes. To determine the fruit density, three medium sized tomatoes per plot were selected randomly. Tomatoes were placed in a box with known volume and weight and then their total weight was measured. The box was filled by a substance with known density (e.g., canola small rounded seed, 0.676 g/cm<sup>3</sup>), that was suitable for filling the empty volume of the box and were weighted again to determine canola's weight. Fruit density obtained from the following equation:

$$\rho_t = \frac{\rho_c - w_t}{(\rho_c \times v_b) - w_c} \quad (4)$$

Where:  $\rho_t$  is the tomato density (gr/cm<sup>3</sup>),  $\rho_c$  is the canola density (gr/cm<sup>3</sup>),  $w_t$  is the tomato weight (gr),  $w_c$  is the canola seed weight (gr) and  $v_b$  is the box volume (cm<sup>3</sup>).

Table 3. Tomato characteristics in different salinities and irrigation management strategies.

Irrigation water (EC) salinity (ds/m)	Plant height (m)	Plant diameter (m)	Water content (%)	Density (g/cm <sup>3</sup> )	L "Factor"	a "Factor"	b "Factor"	Texture strength (load gram)	Sodium concentration (ppm)	Calcium concentration (ppm)	Nitrogen concentration (%)
Irrigation management, m <sub>1</sub>											
S <sub>0</sub> (0.68)	53.33	48.33	96.57	0.94	46.67	57.00	51.67	599.00	7.04	970.33	3.19
S <sub>1</sub> (2)	51.67	50	96.55	0.92	42.67	58.67	49.67	663.83	7.35	891.67	3.06
S <sub>2</sub> (4)	50	48.33	96.38	0.92	40.33	54.67	47.00	742.67	7.38	734.00	2.97
S <sub>3</sub> (6)	48.33	45	96.29	0.94	43.00	58.00	48.67	787.17	9.37	700.00	2.62
S <sub>4</sub> (8)	41.67	38	95.28	0.90	41.00	55.67	47.33	780.00	17.87	673.67	2.61
mean	49 <sup>a</sup>	45.9 <sup>a</sup>	91.46 <sup>a</sup>	0.924 <sup>a</sup>	42.93 <sup>a</sup>	57.73 <sup>a</sup>	48.86 <sup>a</sup>	758.8 <sup>a</sup>	9.8 <sup>a</sup>	793.93 <sup>a</sup>	2.94 <sup>a</sup>
Irrigation management, m <sub>2</sub>											
S <sub>0</sub> (0.68)	53.33	45	96.97	0.90	45.33	58.67	48.67	642.00	6.88	1049.67	3.32
S <sub>1</sub> (2)	51.67	55	96.73	0.91	41.33	56.00	47.67	695.17	8.04	1037.33	3.33
S <sub>2</sub> (4)	51.67	43.33	96.50	0.93	43.67	59.00	50.33	734.00	8.48	916.00	2.74
S <sub>3</sub> (6)	51.67	40	95.98	0.90	42.67	59.00	49.33	834.00	10.24	817.00	2.62
S <sub>4</sub> (8)	51.67	38.33	95.87	0.93	40.00	56.00	47.33	887.50	15.26	667.67	2.68
mean	52 <sup>a</sup>	44.33 <sup>a</sup>	96.35 <sup>a</sup>	0.914 <sup>a</sup>	43.66 <sup>a</sup>	56.8 <sup>a</sup>	48.66 <sup>a</sup>	729.07 <sup>a</sup>	9.78 <sup>a</sup>	897.53 <sup>a</sup>	2.88 <sup>a</sup>
Mean of m <sub>1</sub> and m <sub>2</sub>											
S <sub>0</sub> (0.68)	53.33 <sup>a</sup>	46.66 <sup>ab</sup>	96.77 <sup>a</sup>	0.92 <sup>a</sup>	46 <sup>a</sup>	57.83 <sup>a</sup>	50.17 <sup>a</sup>	620.67 <sup>c</sup>	6.96 <sup>b</sup>	1010 <sup>a</sup>	3.25 <sup>a</sup>
S <sub>1</sub> (2)	51.67 <sup>a</sup>	47.15 <sup>a</sup>	96.64 <sup>ab</sup>	0.915 <sup>a</sup>	42 <sup>ab</sup>	57.33 <sup>a</sup>	48.67 <sup>a</sup>	679.83 <sup>bc</sup>	7.69 <sup>b</sup>	964.5 <sup>a</sup>	3.2 <sup>a</sup>
S <sub>2</sub> (4)	5.83 <sup>a</sup>	45.83 <sup>bc</sup>	96.43 <sup>bc</sup>	0.925 <sup>a</sup>	42.67 <sup>ab</sup>	56.83 <sup>a</sup>	48.66 <sup>a</sup>	729.33 <sup>abc</sup>	7.92 <sup>b</sup>	825 <sup>a</sup>	2.83 <sup>ab</sup>
S <sub>3</sub> (6)	50 <sup>a</sup>	42.50 <sup>bc</sup>	96.14 <sup>bc</sup>	0.92 <sup>a</sup>	42.83 <sup>ab</sup>	58.5 <sup>a</sup>	49 <sup>a</sup>	811 <sup>ab</sup>	9.8 <sup>b</sup>	758.5 <sup>a</sup>	2.65 <sup>b</sup>
S <sub>4</sub> (8)	46.67 <sup>a</sup>	38.17 <sup>c</sup>	95.92 <sup>c</sup>	0.915 <sup>a</sup>	40.5 <sup>c</sup>	55.83 <sup>a</sup>	47.33 <sup>a</sup>	878.83 <sup>a</sup>	16.57 <sup>a</sup>	670.67 <sup>a</sup>	2.62 <sup>b</sup>

Different letters in columns and rows indicate a statistically significant difference at 5% probability.



### *Colorimetric characteristics*

Color and appearance of vegetable, is the first parameter that is considered by the consumers. Human eye has ability in detecting colors, so that it can detect minimal changes in the transparency and light. But this ability is variable in different people. The more conventional way to measure color of food is Lab or L\* a\* b\* method. L\* a\* b\* or CIELab is a global standard that was published by International Lighting Commission where L is luminance, expressed as a percentage (0 for black to 100 for white) and "a" and "b" are two color ranges, from red to green and from yellow to blue, respectively, with values ranging from -120 to +120 (Larrain et al., 2008). To determine the colorimetric characteristics, color photo of tomatoes was taken under the same lighting conditions in a chamber with a white background using high resolution digital camera. Then images were processed by Photoshop software to determine Lab.

### *Tomato firmness*

One of the important characteristics in tomato fruit is good texture and resistance to mechanical damage resulting from transportation. To measure the tomato texture strength TA device (STEVENS-LFRA texture analyzer) was used (Marrs, 1980). Tomato texture strength was measured with the incoming force to crush the fruit.

### *Concentration of sodium and calcium*

To measure the sodium and calcium concentrations in tomato fruit, samples which dried to determine water content were used. The dried samples were powder and passed through the sieve (No. 200) and then ash was made. Five ml HCl per one gram of powder was added to the ash and after passing through filter paper, the volume of sample was brought to 50 cm<sup>3</sup> by boiling distilled water. After preparing plant samples, the sodium and calcium concentrations were measured by Flame Photometer at wavelength of 589 nm and atomic absorption (GBC 932, Perkin Elmer, USA) at wave length of 422.7 nm, respectively (Pauwels et al., 1997).

### Nitrogen concentration

To measure the nitrogen concentration in tomato, fruit samples which dried to determine water content were used. Then samples were powder and passed through the sieve (No. 200). The nitrogen was measured by Kjeldahl method (Bermner and Mulvaney, 1982).

Table 4. Fresh tomato yield (t/ha), in the different irrigation management strategies and salinities.

Irrigation water salinity (EC <sub>i</sub> ) (ds/m)	irrigation management strategies		
	Alternative In season (M <sub>1</sub> )	Alternative In event (M <sub>2</sub> )	mean
	Yeild (t/ha)	Yeild (t/ha)	Yeild (t/ha)
S <sub>0</sub> (0.68)	43.06 <sup>AB*</sup>	32.95 <sup>BCDE</sup>	38.02 <sup>B</sup>
S <sub>1</sub> (2)	54.12 <sup>A</sup>	40.19 <sup>BC</sup>	47.15 <sup>A</sup>
S <sub>2</sub> (4)	27.48 <sup>DEF</sup>	32.30 <sup>BCDE</sup>	31.57 <sup>BC</sup>
S <sub>3</sub> (6)	21.33 <sup>EF</sup>	29.37 <sup>CDE</sup>	25.35 <sup>CD</sup>
S <sub>4</sub> (8)	18.81 <sup>F</sup>	23.58 <sup>DEF</sup>	21.20 <sup>D</sup>
mean	32.97 <sup>A</sup>	31.67 <sup>A</sup>	

\* Mean followed letters in columns and rows are not statistically different, according to LSD test at 5% level of probability.

## Results and Discussion

### Fresh tomato yield

Based on the amount of fresh tomato yield (t/ha) in different treatments (Table 3), analysis of variance was performed. The results showed that the salinity and interaction of salinity and irrigation management strategies on tomato yield, was significant at 5% level of probability. The irrigation management strategies showed no significant effect on tomato yield. The maximum and minimum of yield were obtained in M<sub>1</sub>S<sub>1</sub> (54.12 t/ha) and M<sub>1</sub>S<sub>4</sub> (18.81 t/ha) treatments, respectively. Fresh tomato yield in M<sub>1</sub>S<sub>1</sub> and M<sub>2</sub>S<sub>1</sub> treatments were 25.7% and 22.0% more than the control salinity level (S<sub>0</sub>), respectively. According to Table 3 and Figure 2, maximum yield in both irrigation management strategies were in S<sub>1</sub> salinity level (2 dS/m). In S<sub>1</sub>, S<sub>2</sub>, S<sub>3</sub> and S<sub>4</sub> salinity levels, the changes of yields relative to control treatment (S<sub>0</sub>) were +25.7, -36.2, -50.5, for M<sub>1</sub> treatments and -56.3%; +22.0, -2.0, -10.9, -28.4%; for M<sub>2</sub> treatment and +23.8, -19.1, -30.7, -42.4%, for mean of M<sub>1</sub> and M<sub>2</sub> treatments, respectively.




Figure 2. Relationship between Fresh tomato yield and salinity levels of irrigation water at mean of  $M_1$  and  $M_2$  treatments.

#### *Yield salinity relationship*

Relationship between fresh tomato yield and water salinity at two irrigation management strategies are as follow:

For mean of alternate in season ( $M_1$ ) and alternate in event ( $M_2$ ) treatments:

$$y = -0.196EC^4 + 3.669EC^3 - 22.81EC^2 + 49.05EC + 14.089 \quad (5)$$

$$n = 30 \quad R^2 = 0.69 \quad Se = 6.78 \quad P - value = 3.93 \times 10^{-6}$$

*For alternate in season ( $M_1$ ) treatments*

$$y = -0.281EC^4 + 5.279EC^3 - 32.644EC^2 + 68.491EC + 9.980 \quad (6)$$

$$n = 15 \quad R^2 = 0.81 \quad Se = 6.43 \quad P - value = 2.07 \times 10^{-4}$$

*For alternate in event ( $M_2$ ) treatments*

$$y = -0.111EC^4 + 2.059EC^3 - 12.980EC^2 + 29.6008EC + 18.198 \quad (7)$$

$$n = 15 \quad R^2 = 0.77 \quad Se = 3.56 \quad P - value = 2.83 \times 10^{-3}$$

Where:  $y$  is the fresh yield of tomato (t/ha) and  $EC_i$  is the electrical conductivity of irrigation water (dS/m). In Eq. (5), until the salinity of 2 dS/m, yield was increased and then with increasing salinity, yield was decreased. The trend of yield reduction is high until the salinity reached-5 dS/m and then it become low-at salinity of 8 dS/m. According to Figure 2, for mean of  $M_1$  and  $M_2$  strategies,  $EC_i$  values that reduce 10, 25, 50, 75 and 100% of yield relative to control were 2.4, 3.3, 5.2, 8.5 and 8.92 dS/m, respectively.

### Water Productivity

Water productivity (WP) was obtained from division of the fresh tomato yield (kg/ha) by amount of irrigation water applied to the farm during the growing season (mm). The plant yield (kg/ha) and water productivity for different treatments are shown in Table 5. The amounts of applied water with increasing salinity levels were reduced. Therefore, with increased salinity, WP increased. In general, in  $M_1$  treatment WP was higher than  $M_2$  treatment. Therefore, the WP of  $M_1S_0$ ,  $M_1S_1$  and  $M_1S_4$  treatments were 30.7, 49.4 and 9.6% higher than  $M_2S_0$ ,  $M_2S_1$  and  $M_2S_4$  treatments, respectively. However, in  $M_1S_2$  and  $M_1S_3$  treatments WP were 11.8 and 12.3% lower than  $M_2S_2$  and  $M_2S_3$  treatments, respectively. Therefore,  $M_1$  treatment in low ( $S_0$ ,  $S_1$ ) and high ( $S_4$ ) salinity levels, showed higher WP. However, in medium salinity levels ( $S_2$ ,  $S_3$ ),  $M_1$  treatment resulted in lower WP than  $M_2$  treatment.

Table 5. Yield and water productivity in different treatments.

Water Productivity (WP) (kg/m <sup>3</sup> )	Applied water (mm)	$EC_i$ (dS/m)	Management Strategy
3.417	1260	0.68	$S_0$
5.306	1020	2	$S_1$
3.523	780	4	$S_2$
3.949	540	6	$S_3$
6.272	300	8	$S_4$
2.615	1260	0.68	$S_0$
3.550	1132	2	$S_1$
3.995	892	4	$S_2$
4.504	652	6	$S_3$
5.722	412	8	$S_4$

Soil salinity ( $EC_e$ )

Figure 3 shows the values of soil saturated extract ( $EC_e$ ) at the end of growing season for each treatment at different soil depths. In all soil layers, the lowest  $EC_e$  was in  $M_1S_0$  treatment. Highest  $EC_e$  in all soil layers was with  $S_4$  level. Further, in the first (0-30 cm) and second layers (30-50 cm)  $EC_e$  was highest in  $M_2$ , however, in third (50-70 cm) and fourth layers (70-90 cm) it was highest in  $M_1$  treatment. In salinity of 0.68 and 2 dS/m,  $M_1$  treatment and in salinity levels over 2 dS/m,  $M_2$  treatments have lower mean  $EC_e$ . Similar to WP, for soil salinity the  $M_2$  treatment is appropriate management at high water salinity levels.




Figure 3.  $EC_e$  in different soil depths and management strategies: a)  $M_1$  strategy, b)  $M_2$  strategy.

According to Figure 3, the use of saline water to irrigate crops in all growing season increased  $EC_e$  for both management strategies. For salinity between 0.68-6 dS/m,  $EC_e$  in  $M_2$  treatment were close to each other, however, for salinity of 8 dS/m, the  $EC_e$  increased considerably. The  $EC_e$  in  $M_2$  treatment for all salinity levels is less than  $M_1$  treatment due to more effective leaching in  $M_2$  management strategy. Salts were accumulated in 0-50 cm layer and the  $EC_e$  was decreased in depth lower than 50 cm due to the presence of a layer with low hydraulic conductivity in this depth. In general,  $M_2$  management strategy, due to more efficient leaching in the upper layers, is more appropriate than  $M_1$  for tomato plant and may be for growing shallow root crops.

#### *Plant height and canopy diameter*

Statistical results were showed that salinity, irrigation management strategies and the interaction of salinity and irrigation management strategies has no significant effect on plant height at 5% level. The maximum of plant height was in  $M_1S_0$  and  $M_2S_0$  (53.3 cm) and the minimum was in  $M_1S_4$  (41.7 cm) treatments, respectively. For canopy diameter, the results showed that salinity has a significant effect at 5% level while the irrigation management strategies and the interaction of salinity and irrigation management strategies have no significant effect ( $P < 0.05$ ). The maximum and minimum of canopy diameter were in  $M_2S_1$  (55 cm) and  $M_1S_4$  (38 cm) treatments, respectively. Similar to behavior of yield response to salinity levels, the maximum canopy diameter in both management strategies was in  $S_1$  salinity level and for salinity over 2 dS/m; canopy diameter was reduced with increasing salinity level. In general, the osmotic pressure of soil solution increases. Therefore, the amount of energy that plant spent to absorb water increases with increasing salinity level. This is why, the yield and canopy diameter were reduced with increasing salinity.

#### *Tomato water content*

Based on analysis of variance, salinity showed a significant effect while the irrigation management strategies and the interaction of salinity and irrigation management strategies have no significant effect on tomato water content at 5% level of probability. The maximum and minimum values of tomato water content were in  $M_2S_0$  (97.0%) and  $M_2S_4$  (95.9%) treatments,

respectively. Tomato water content reduced with increasing the salinity level. The average of tomato water content in M<sub>1</sub> management strategy at low and medium salinity levels (0.68, 2 and 4 dS/m) was lower than M<sub>2</sub> treatment, but at high salinity levels (6 and 8 dS/m) was higher than M<sub>2</sub> treatment. According to these results, the S<sub>4</sub> salinity level and M<sub>1</sub> management strategy was increased the tomato water content.

#### *Fruit density*

Statistical results showed that salinity, irrigation management strategies and the interaction of salinity and irrigation management strategies have no significant effect on fruit density at 5% level of probability. Therefore, changes in fruit density, has not a specific trend with respect to salinities and management strategies. The maximum and minimum values of fruit density were in M<sub>1</sub>S<sub>3</sub> (0.94 g/cm<sup>3</sup>) and M<sub>1</sub>S<sub>4</sub> (0.90 g/cm<sup>3</sup>) treatments, respectively.

#### *Colorimetric characteristics*

Based on analysis of variance, salinity level has a significant effect while the irrigation management strategies and the interaction of salinity and irrigation management strategies have no significant effect on "L" factor at 5% level of probability. The maximum and minimum values of "L" factor were in M<sub>1</sub>S<sub>0</sub> (46.7) and M<sub>2</sub>S<sub>4</sub> (40.0) treatments, respectively. The "L" factor reduced with increasing the salinity level.

The interaction of salinity and irrigation management strategies has a significant effect while the salinity and irrigation management strategies have no significant effect on "a" factor at 5% level of probability. The maximum value of "a" factor was in M<sub>1</sub>S<sub>0</sub> and M<sub>2</sub>S<sub>0</sub> (59.0) and the minimum was in M<sub>1</sub>S<sub>4</sub> (54.7) treatments, respectively. Higher value of "L" and "a" factors represent premature fruit (Larrain et al., 2008).

There is not a trend between salinity, irrigation management strategies and interaction of salinity and irrigation management on "b" factor. The maximum and minimum values of "b" factor were in M<sub>1</sub>S<sub>0</sub> (51.7) and M<sub>1</sub>S<sub>2</sub> (47.0) treatments, respectively. The M<sub>1</sub>S<sub>2</sub> was also the optimum treatment according to "L" and "a" factors. Therefore, salinity has significant effect on the just "L" factor. The irrigation management strategies and interaction of salinity and irrigation management have not significant effect on the "L", "a" and "b" factors. According to these results the S<sub>4</sub> salinity level has the

positive effect on the "L" and "a" factors. In general, the tomato fruits that were irrigated with saline water had been riper. Therefore, use of saline water can be considered the optimal method to precocity of fruits.

#### *Tomato firmness*

Based on analysis of variance, salinity showed a significant effect while the irrigation management strategies and the interaction of salinity and irrigation management strategies have no significant effect on tomato texture strength at 5% level of probability. The maximum and minimum values of tomato texture strength were in  $M_2S_4$  (887.5 load gram) and  $M_1S_0$  (559 load gram) treatments, respectively. In general, the tomato texture strength increased with increasing salinity levels. In  $S_1$ ,  $S_2$ ,  $S_3$  and  $S_4$  levels the tomato texture strength was 10, 18, 31 and 42% higher than the control treatments, respectively. These results were in agreement with results obtained on tomato water content. The tomato water content was reduced and then the texture strength was increased with increasing salinity levels. According to these results the  $S_4$  salinity level has the positive effect on the texture strength.

#### *Sodium, calcium and nitrogen concentration*

Based on analysis of variance, salinity showed a significant effect just on sodium and nitrogen concentration while the salinity for calcium and irrigation management strategies and the interaction of salinity and irrigation management strategies have no significant effect at 1% level of probability on sodium, calcium and nitrogen concentration. Sodium concentration increased with increasing salinity levels. For  $M_1$  and  $M_2$  treatments, the sodium concentrations in  $S_1$ ,  $S_2$ ,  $S_3$  and  $S_4$  salinity levels were increased 4.4, 4.8, 33.1, 153.8% and 16.9, 23.2, 48.8, 121.8% relative to control ( $S_0$ ), respectively. Therefore, increasing sodium concentration in  $S_1$ ,  $S_2$  and  $S_3$  salinity levels in  $M_1$  treatment was lower than those in  $M_2$  treatment. However, it was inverse in  $S_4$  salinity level. The maximum and minimum values of sodium concentrations were in  $M_1S_4$  (17.9  $\mu\text{g/g}$ ) and  $M_2S_0$  (6.88  $\mu\text{g/g}$ ) treatments, respectively.

The nitrogen and calcium concentrations were decreased with increasing salinity levels. In  $M_1$  and  $M_2$  treatments, calcium concentration in  $S_1$ ,  $S_2$ ,  $S_3$  and  $S_4$  salinity levels were decreased 8.1, 24.4, 27.9, 30.6% and 1.2, 12.4, 22.2, 36.4% relative to control ( $S_0$ ), respectively. The maximum and minimum values of nitrogen concentration were in  $M_2S_1$  (3.3%) and  $M_1S_4$



(2.6%) treatments, respectively. In saline conditions, increased concentrations of salts prevent the natural nutrition in plants. Nitrogen is one of the most important elements that limiting plant growth. Nitrogen uptake is reduced more than any other nutrient elements when plants are under salt stress. In saline conditions, many factors are effective in the reduction of nitrogen uptake by plants such as reduction of the permeability of plant roots, reduction of the microbial activity and subsequent loss of soil mineralization of organic compounds, high concentrations of chloride in the root zone and reduction of the nitrification. According to these results the  $S_4$  and  $S_1$  salinity levels and  $M_1$  management strategy have the positive effect on the sodium and nitrogen concentrations, respectively.

### **Conclusion**

With increasing salinity levels, canopy diameter, fruit water content, "a" and "L" factors of colorimetric characteristics and nitrogen concentration were decreased. However, sodium concentration and texture strength were increased. Plant height, "b" factor in the colorimetric characteristics and calcium concentration, were not influenced by salinity levels. Highest and lowest fresh yield were obtained in  $M_1S_1$  (54.12 t/ha) and  $M_1S_4$  treatments (18.81 t/ha), respectively. Tomato fresh yields in  $M_1S_1$  and  $M_2S_1$  treatments were 25.7% and 22.0%, higher than the yield in the control salinity, respectively. Therefore, in both irrigation management strategies, in  $S_1$  salinity level (2 dS/m), yield was higher than  $S_0$  (0.68 dS/m). Water productivity increased with increasing salinity, so that highest water productivity was obtained in  $M_1S_4$  treatment. However, the acceptable WP with good yield was obtained in  $M_1S_1$  treatment. The  $M_1$  management strategy in low ( $S_0$ ,  $S_1$ ) and high ( $S_4$ ) salinity levels showed higher WP. However, medium salinity levels ( $S_2$ ,  $S_3$ ) have lower WP than  $M_2$  treatment. In terms of impact on soil salinity and crop yield similar to WP, at lower levels of salinity (0.68 and 2 dS/m) the  $M_1$  management strategy and at salinity levels over 4 dS/m, the  $M_2$  strategy is an appropriate management strategy. In general,  $M_1$  treatment has better results than  $M_2$  treatment.

### **Acknowledgement**

The authors wish to thank the Research Council of Shiraz University and National Drought Research Institute of Iran for financial support.

## References

- Anonymous, 2009. Sustainable management of unconventional water resources and saline soils. National Research centre for salinity, Yazd, Iran.
- Borg, H., Grimes, D.W., 1986. Depth development of roots with time: An empirical description. *Trans: ASAE*, 29 (1), 194-197.
- Bermner, J.M., Mulvaney, C.S., 1982. Total nitrogen, pp. 595-624. In: A.L. Page (ed) *Methods of soil Analysis. Part II. 2<sup>nd</sup> Ed.*
- Holland, B., Welch, A., Unwin, I., 1991. *The Composition of Foods*. Royal Society of Chemistry, Cambridge, pp. 268-271.
- Larrain, R.E., Schaefer, D.M., Reed, J.D., 2008. Use of digital images to estimate CIE color coordinates of beef. *Food Research International*, 41, 380-385.
- Malash, N., Flowers, T.J., Ragab, R., 2005. Effect of irrigation systems and water management practices using saline and non-saline water on tomato production. *Agricultural Water Management*, 78, 25-38.
- Malash, N., Ali, F.A., Fatahalla, M.A., 2008. Response of tomato to irrigation with saline water applied by different irrigation methods and water management strategies. *Inter. J. Plant Prod.* 2 (2), 1735-8043.
- Marrs, W.M., 1980. The Stevens-LFRA texture analyzer: a new instrument for gel testing and for the evaluation of texture. *The British Food Manufacturing Industries Research Assoc.* 25p.
- Murtaza, G., Ghafoor, A., Qadir, M., 2006. Irrigation and soil management strategies for using saline-sodic water in a cotton-wheat rotation. *Agricultural Water Management*, 81, 98-114.
- Narash, R.K., Minhas, P.S., Goyal, A.K., Chauhan, C.P.S., Gupta, P.K., 1993. Conjunctive use of saline and non-saline waters. II. Field comparison of cyclic uses and mixing for wheat. *Agricultural Water Management*, 23, 139-148.
- Pasternak, D., De Malach, Y., Borovic, I., 1986. Irrigation with brackish water under desert condition VII. Effect of time and application of brackish water on production of processing tomatoes (*lycopesionesculentum* Mill.). *Agricultural Water Management*, 12, 149-158.
- Pasternak, D., De Malach, Y., 1995. Irrigation with brackish water under desert condition X. Irrigation management of tomatoes (*lycopesionesculentum* Mill.) on desert sand dunes. *Agricultural Water Management*, 28, 121-132.
- Pauwels, P.J., Tardif, S., Wurch, T., Colpaert, F., 1997. Stimulated [35S]GTP $\gamma$ S binding by 5HT1A receptor agonists in recombinant cell lines. *Naunyn Schmiedebergs Arch. Pharmacol.* 356, 551-561.
- Wallace, J.S., 2000. Increasing agricultural water use efficiency to meet future food production. *Agricultural Water Management*, 82, 105-119.
- Wan, Sh., Kang, Y., Wang, D., Liu, Sh.P., 2010. Effect of saline water on cucumber (*Cucumissativus* L.) yield and water use under drip irrigation on North China. *Agricultural Water Management*, 98 (1), 105-113.